

The listing of claims will replace all prior versions and listing of claims in the application:

**Listing of Claims:**

Claim 1. (original) A water-dispersible, freeze-dried bioavailable coenzyme Q-10/cyclodextrin complex.

Claim 2. (original) The complex of claim 1, wherein the molar ratio of cyclodextrin to coenzyme Q-10 ranges from about 0.5:1 to 10:1.

Claim 3. (original) The complex of claim 2, wherein said molar ratio ranges from about 1:1 to 2:1.

Claim 4. (original) The complex of claim 1, wherein said cyclodextrin is one or more of  $\beta$ -cyclodextrin or  $\gamma$ -cyclodextrin.

Claim 5. (currently amended) The complex of claim 1, which formulated into one or more of a topical preparation, a sublingual formulation, or for oral ingestion.

Claim 6. (currently amended) A method for making a water-dispersible complex, which comprises the steps of:

- (a) preparing an aqueous slurry of a coenzyme Q-10/cyclodextrin complex by adding coenzyme Q-10 to an aqueous dispersion of cyclodextrin; and
- (b) drying by one or more of spray drying, vacuum-drying, or freeze drying, said aqueous slurry to produce said complex.

Claim 7. (original) The method of claim 6, wherein the molar ratio of cyclodextrin to coenzyme Q-10 ranges from about 0.5:1 to 10:1.

Claim 8. (original) The method of claim 7, wherein said molar ratio ranges from about 1:1 to 2:1.

Claim 9. (original) The method of claim 6, wherein said cyclodextrin is one or more of  $\beta$ -cyclodextrin or  $\gamma$ -cyclodextrin.

Claim 10. (currently amended) A method for ~~treating~~ administering to an animal with a bioavailable coenzyme Q-10 complex, which comprises the steps of:

- (a) preparing a water-dispersible coenzyme Q-10/cyclodextrin complex by adding coenzyme Q-10 to an aqueous dispersion of cyclodextrin; and
- (b) administering said complex to an animal.

Claim 11. (original) The method of claim 10, wherein said animal is a human.

Claim 12. (original) The method of claim 10, wherein said complex is ingested by said animal.

Claim 13. (original) The method of claim 10, wherein the molar ratio of cyclodextrin to coenzyme Q-10 ranges from about 0.5:1 to 10:1.

Claim 14. (original) The method of claim 13, wherein said molar ratio ranges from about 1:1 to 2:1.

Claim 15. (original) The method of claim 10, wherein said cyclodextrin is one or more of  $\beta$ -cyclodextrin or  $\gamma$ -cyclodextrin.

Claim 16. (original) The method of claim 10, wherein said complex is prepared by freeze-drying.

Cancel Claim 17.

Cancel Claim 18.

Claim 19. (original) The method of claim 10, which formulated into one or more of a topical preparation, a sublingual formulation, or for oral ingestion.

Claim 20. (original) The method of claim 17, wherein said cyclodextrin is one or more of  $\beta$ -cyclodextrin or  $\gamma$ -cyclodextrin.

Claim 21. (New)        The method of claim 6, wherein said adding coenzyme Q-10 to an aqueous dispersion of cyclodextrin in step (a) is accomplished at about room temperature using homogenization followed by storage under refrigeration.

Claim 22. (New)        The method of claim 10, wherein said adding coenzyme Q-10 to an aqueous dispersion of cyclodextrin in step (a) is accomplished at about room temperature using homogenization followed by storage under refrigeration.